arXiv:cond-mat/0605128v1 [cond-mat.stat-mech] 5 May 2006

Exact expressions of mean first-passage times and splitting probabilities for random walks in bounded rectangular domains

S. Condamin and O. Bénichou

Université Pierre et Marie Curie-Paris6,

Laboratoire de Physique Théorique de la Matière Condensée,

UMR-CNRS 7600, case courrier 121, 4 Place Jussieu, F-75005 Paris

(Dated: February 4, 2008)

PACS numbers: 05.40Fb, 05.40Jc

Recently, we have proposed a novel computation method of first-passage times of a random walker between a starting site and a target site of regular bounded lattices of arbitrary shape[1]. Such first passage time properties are for example crucial to describe the kinetics of diffusion limited reactions in confined media[2, 3]. The obtained expressions involve pseudo-Green functions[4], which have been estimated according to different approximation schemes. In this note, we give the exact expression of these pseudo-Green functions in both cases of a rectangular domain with reflecting boundaries and a rectangular domain with periodic boundary conditions. This allows us to provide exact and explicit expressions of mean first-passage times with one or two target sites.

Consider first a random walker starting at the source S, of position $\mathbf{r}_S$ of a rectangular regular lattice with reflecting boundary conditions. The mean time it takes to reach the target T, of position $\mathbf{r}_T$, for the first time, is given by[1]

$$\langle \mathbf{T} \rangle = N[H(\mathbf{r}_T|\mathbf{r}_T) - H(\mathbf{r}_T|\mathbf{r}_S)] \tag{1}$$

where H is the pseudo-Green function, which satisfies

$$H(\mathbf{r}_i|\mathbf{r}_j) = \frac{1}{\sigma}\sum_{\langle k,i \rangle} H(\mathbf{r}_k|\mathbf{r}_j) + \delta_{ij} - \frac{1}{N}, \tag{2}$$

σ being the coordination number of the lattice, namely 4 for a 2-D lattice, or 6 for a 3-D lattice. In this expression, the sum runs over all the neighbours k of the site i, considering that a site which is at the boundary of the domain is its own neighbour. N is the total number of sites of the lattice. In the presence of two absorbing targets, the mean time it takes to reach either of the two targets is:

$$\langle \mathbf{T} \rangle = N\frac{(H_{01} - H_{1s})(H_{02} - H_{2s}) - (H_{12} - H_{2s})(H_{12} - H_{1s})}{H_{01} + H_{02} - 2H_{12}} \tag{3}$$

while the eventual hitting probabilities P_i to reach the target i writes

$$\begin{cases} P_1 = \frac{H_{1s}+H_{02}-H_{2s}-H_{12}}{H_{01}+H_{02}-2H_{12}} \\ P_2 = \frac{H_{2s}+H_{01}-H_{1s}-H_{12}}{H_{01}+H_{02}-2H_{12}} \end{cases} \tag{4}$$

where $H_{12} = H(\mathbf{r}_{T_1}|\mathbf{r}_{T_2})$ and, for $i = 1$ or 2, $H_{is} = H(\mathbf{r}_{T_i}|\mathbf{r}_S)$, $H_{0i} = H(\mathbf{r}_{T_i}|\mathbf{r}_{T_i})$.

The exact expression of the pseudo-Green function H involved in previous equations may be computed explicitely with the help of Fourier analysis. For a 2D domain with X sites in



FIG. 1: 2D Two-target simulations (red crosses) vs. theory (plain line). One target is fixed at (-5,0); The source is fixed at (5,0); The other target is at (x,3). The domain is a square of side 41, the middle is the point (0,0). The absorption time is normalized by the number of sites N.

the x direction and Y sites in the y direction, H writes

$$\begin{aligned} H(\mathbf{r}|\mathbf{r}') \;=\; & \frac{1}{N}\sum_{m=1}^{X-1}\sum_{n=1}^{Y-1}\frac{4\cos\frac{m\pi x'}{X}\cos\frac{n\pi y'}{Y}\cos\frac{m\pi x}{X}\cos\frac{n\pi y}{Y}}{1-\frac{1}{2}\left(\cos\frac{m\pi}{X}+\cos\frac{n\pi}{Y}\right)} \\ & +\frac{1}{N}\sum_{m=1}^{X-1}\frac{4\cos\frac{m\pi x'}{X}\cos\frac{m\pi x}{X}}{1-\cos\frac{m\pi}{X}} \\ & +\frac{1}{N}\sum_{n=1}^{Y-1}\frac{4\cos\frac{n\pi y'}{Y}\cos\frac{n\pi y}{Y}}{1-\cos\frac{n\pi}{Y}} \end{aligned} \tag{5}$$

where x and y are the coordinates of $\mathbf{r}$ and x' and y' those of $\mathbf{r}'$. Here, the coordinates of the left-bottom corner are taken equal to $(1/2, 1/2)$, so that all the coordinates are half-integers.

Figure 1 shows the mean absorption time $\langle\mathbf{T}\rangle$ given by eq. (3) as a function of the target positions together with numerical simulations in the case of a 2D square domain. These simulations have been performed with a method based on the exact enumeration method[5].

Similar results can be obtained in the case of a rectangular domain with periodic boundary conditions. Here, the pseudo-Green function is

$$H(\mathbf{r}|\mathbf{r}') = \frac{1}{N}\sum_{m=0}^{X-1}\sum_{n=\delta_{0m}}^{Y-1}\frac{\exp\frac{2im\pi(x-x')}{X}\exp\frac{2ni\pi(y-y')}{Y}}{1-\frac{1}{2}\left(\cos\frac{2m\pi}{X}+\cos\frac{2n\pi}{Y}\right)} \tag{6}$$

where δ_{0m} is the Kronecker delta.

Note that our results can be easily extended to the 3D case.

We gratefully thank S. Redner for suggesting us the simulation method.

---

[1] S. Condamin, O. Bénichou and M. Moreau, Phys. Rev. Lett. **95**, 260601 (2005).

[2] S. Rice, *Diffusion-Limited Reactions* (Elsevier, Amsterdam, 1985).

[3] S. Redner, *A guide to First- Passage Processes* (Cambridge University Press, Cambridge, 2001)

[4] G. Barton, *Elements of Green's Functions and Propagation* (Oxford Science Publications, Oxford, 1989)

[5] I. Majid, D. B. Avraham, S. Havlin, and H. E. Stanley, Phys. Rev. B **30**, 1626 (1984).